

Precision Measurements of the Mass, the Widths of $\psi(3770)$ Resonance and the Cross Section $\sigma[e^+e^- \rightarrow \psi(3770)]$ at $E_{\text{cm}} = 3.7724$ GeV

M. Ablikim¹, J. Z. Bai¹, Y. Ban¹¹, J. G. Bian¹, X. Cai¹, H. F. Chen¹⁵, H. S. Chen¹, H. X. Chen¹, J. C. Chen¹, Jin Chen¹, Y. B. Chen¹, S. P. Chi², Y. P. Chu¹, X. Z. Cui¹, Y. S. Dai¹⁷, Z. Y. Deng¹, L. Y. Dong^{1a}, Q. F. Dong¹⁴, S. X. Du¹, Z. Z. Du¹, J. Fang¹, S. S. Fang², C. D. Fu¹, C. S. Gao¹, Y. N. Gao¹⁴, S. D. Gu¹, Y. T. Gu⁴, Y. N. Guo¹, Y. Q. Guo¹, K. L. He¹, M. He¹², Y. K. Heng¹, H. M. Hu¹, T. Hu¹, X. P. Huang¹, X. T. Huang¹², X. B. Ji¹, X. S. Jiang¹, J. B. Jiao¹², D. P. Jin¹, S. Jin¹, Yi Jin¹, Y. F. Lai¹, G. Li², H. B. Li¹, H. H. Li¹, J. Li¹, R. Y. Li¹, S. M. Li¹, W. D. Li¹, W. G. Li¹, X. L. Li⁸, X. Q. Li¹⁰, Y. L. Li⁴, Y. F. Liang¹³, H. B. Liao⁶, C. X. Liu¹, F. Liu⁶, Fang Liu¹⁵, H. H. Liu¹, H. M. Liu¹, J. Liu¹¹, J. B. Liu¹, J. P. Liu¹⁶, R. G. Liu¹, Z. A. Liu¹, F. Lu¹, G. R. Lu⁵, H. J. Lu¹⁵, J. G. Lu¹, C. L. Luo⁹, F. C. Ma⁸, H. L. Ma¹, L. L. Ma¹, Q. M. Ma¹, X. B. Ma⁵, Z. P. Mao¹, X. H. Mo¹, J. Nie¹, H. P. Peng¹⁵, N. D. Qi¹, H. Qin⁹, J. F. Qiu¹, Z. Y. Ren¹, G. Rong¹, L. Y. Shan¹, L. Shang¹, D. L. Shen¹, X. Y. Shen¹, H. Y. Sheng¹, F. Shi¹, X. Shi^{11b}, H. S. Sun¹, J. F. Sun¹, S. S. Sun¹, Y. Z. Sun¹, Z. J. Sun¹, Z. Q. Tan⁴, X. Tang¹, Y. R. Tian¹⁴, G. L. Tong¹, D. Y. Wang¹, L. Wang¹, L. S. Wang¹, M. Wang¹, P. Wang¹, P. L. Wang¹, W. F. Wang^{1c}, Y. F. Wang¹, Z. Wang¹, Z. Y. Wang¹, Zhe Wang¹, Zheng Wang², C. L. Wei¹, D. H. Wei¹, N. Wu¹, X. M. Xia¹, X. X. Xie¹, B. Xin^{8d}, G. F. Xu¹, Y. Xu¹⁰, M. L. Yan¹⁵, F. Yang¹⁰, H. X. Yang¹, J. Yang¹⁵, Y. X. Yang³, M. H. Ye², Y. X. Ye¹⁵, Z. Y. Yi¹, G. W. Yu¹, C. Z. Yuan¹, J. M. Yuan¹, Y. Yuan¹, S. L. Zang¹, Y. Zeng⁷, Yu Zeng¹, B. X. Zhang¹, B. Y. Zhang¹, C. C. Zhang¹, D. H. Zhang¹, H. Y. Zhang¹, J. W. Zhang¹, J. Y. Zhang¹, Q. J. Zhang¹, X. M. Zhang¹, X. Y. Zhang¹², Yiyun Zhang¹³, Z. P. Zhang¹⁵, Z. Q. Zhang⁵, D. X. Zhao¹, J. W. Zhao¹, M. G. Zhao¹, P. P. Zhao¹, W. R. Zhao¹, H. Q. Zheng¹¹, J. P. Zheng¹, Z. P. Zheng¹, L. Zhou¹, N. F. Zhou¹, K. J. Zhu¹, Q. M. Zhu¹, Y. C. Zhu¹, Y. S. Zhu¹, Yingchun Zhu^{1e}, Z. A. Zhu¹, B. A. Zhuang¹, X. A. Zhuang¹, B. S. Zou¹

(BES Collaboration)

¹ Institute of High Energy Physics, Beijing 100049, People's Republic of China

² China Center for Advanced Science and Technology(CCAST), Beijing 100080, People's Republic of China

³ Guangxi Normal University, Guilin 541004, People's Republic of China

⁴ Guangxi University, Nanning 530004, People's Republic of China

⁵ Henan Normal University, Xinxiang 453002, People's Republic of China

⁶ Huazhong Normal University, Wuhan 430079, People's Republic of China

⁷ Hunan University, Changsha 410082, People's Republic of China

⁸ Liaoning University, Shenyang 110036, People's Republic of China

⁹ Nanjing Normal University, Nanjing 210097, People's Republic of China

¹⁰ Nankai University, Tianjin 300071, People's Republic of China

¹¹ Peking University, Beijing 100871, People's Republic of China

¹² Shandong University, Jinan 250100, People's Republic of China

¹³ Sichuan University, Chengdu 610064, People's Republic of China

¹⁴ Tsinghua University, Beijing 100084, People's Republic of China

¹⁵ University of Science and Technology of China, Hefei 230026, People's Republic of China

¹⁶ Wuhan University, Wuhan 430072, People's Republic of China

¹⁷ Zhejiang University, Hangzhou 310028, People's Republic of China

^a Current address: Iowa State University, Ames, IA 50011-3160, USA

^b Current address: Cornell University, Ithaca, NY 14853, USA

^c Current address: Laboratoire de l'Accélérateur Linéaire, Orsay, F-91898, France

^d Current address: Purdue University, West Lafayette, IN 47907, USA

^e Current address: DESY, D-22607, Hamburg, Germany

By analyzing the R values measured at 68 energy points in the energy region between 3.650 and 3.872 GeV reported in our previous paper, we have precisely measured the mass, the total width, the leptonic width and the leptonic decay branching fraction of the $\psi(3770)$ to be $M_{\psi(3770)} = 3772.4 \pm 0.4 \pm 0.3$ MeV, $\Gamma_{\psi(3770)}^{\text{tot}} = 28.6 \pm 1.2 \pm 0.2$ MeV, $\Gamma_{\psi(3770)}^{ee} = 279 \pm 11 \pm 13$ eV and $B[\psi(3770) \rightarrow e^+e^-] = (0.98 \pm 0.04 \pm 0.04) \times 10^{-5}$, respectively, which result in the observed cross section $\sigma^{\text{obs}}[e^+e^- \rightarrow \psi(3770)] = 7.25 \pm 0.27 \pm 0.34$ nb at $\sqrt{s} = 3772.4$ MeV. We have also measured $R_{\text{uds}} = 2.121 \pm 0.023 \pm 0.084$ for the continuum light hadron production in the region from 3.650 to 3.872 GeV.

PACS numbers: 13.85.Lg, 12.38.Qk, 14.40.Gx, 14.40.Lb

Precise measurements of the mass $M_{\psi(3770)}$, the total width $\Gamma_{\psi(3770)}^{\text{tot}}$ and the leptonic width $\Gamma_{\psi(3770)}^{ee}$ of

the $\psi(3770)$ resonance, and the cross section $\sigma[e^+e^- \rightarrow \psi(3770)]$ for the $\psi(3770)$ production at $\sqrt{s} = M_{\psi(3770)}$

in the e^+e^- annihilation are important in understanding the nature of the $\psi(3770)$. The $\psi(3770)$ is thought to be a mixture of the D -wave and S -wave of the angular momentum eigenstates of $c\bar{c}$ system. The detailed mixing scheme affects the $\psi(3770)$ production and decays. So precise measurements of these quantities would give us some useful information about the nature of the $\psi(3770)$ and offer some insights into the internal wave functions of the charmonium, which are beneficial to understand the dynamics of the 1^{--} resonance production in the e^+e^- annihilation. Potential or quarkonium models based on QCD can calculate the masses [1, 2, 3] of the $c\bar{c}$ bound states, the total widths of the $\psi(3770)$ and other resonances [2, 3], and the width for $\psi(3770) \rightarrow D\bar{D}$ [3, 4]. The Lattice QCD (LQCD) can calculate the mass spectra of the $Q\bar{Q}$ system (heavy quark and anti-quark system, such as the $c\bar{c}$ and $b\bar{b}$ system) [5]. If one can more precisely measure the masses and the widths of the $Q\bar{Q}$ system, the results of these measurements can be used to test the calculation of the quantities by the models and by the LQCD theory. Moreover, these measurements can in turn be used to extract two fundamental parameters in QCD, the c -quark mass and the strong coupling constant $\alpha_s(s)$ at this mass scale [5].

In recent days, new results of the measured quantities related to the $\psi(3770)$ production and decays were reported [6, 7]. These improve our knowledge on charmonium production and decays, especially improve our understanding of the nature of the $\psi(3770)$ resonance. In our previous work [9] we have reported measurements of the $R(s)$ values measured at 68 energy points in the region between 3.650 and 3.872 GeV, where the quantity $R(s)$ is defined as the ratio $\sigma(e^+e^- \rightarrow \text{hadrons})/\sigma(e^+e^- \rightarrow \mu^+\mu^-)$, with $\sigma(e^+e^- \rightarrow \mu^+\mu^-) = 3\pi\alpha^2(0)/4s$, here s is the c.m. (center-of-mass) energy squared and $\alpha(0)$ is the fine structure constant at the lowest energy limit. In this Letter we report the results obtained by further analyzing these $R(s)$ values. From this analysis we obtain $M_{\psi(3770)}$, $\Gamma_{\psi(3770)}^{\text{tot}}$, $\Gamma_{\psi(3770)}^{ee}$ and the leptonic branching fraction for $\psi(3770) \rightarrow e^+e^-$ with improved precision compared to those of the PDG [8] world average, and we obtain the cross section $\sigma[e^+e^- \rightarrow \psi(3770) \rightarrow \text{hadrons}]$ at $\sqrt{s} = 3772.4$ MeV with a precision better than any of those measured in cross section scan experiments previously.

In Ref. [9] we reported measurements of the quantities R_{uds} , $R_{\text{had}}(s)$ and $R_{\text{uds}(c)+\psi(3770)}$ which are the $R(s)$ values for the continuum light hadron production around the $D\bar{D}$ threshold, $R(s)$ values including the contributions from the continuum hadrons and all 1^{--} resonances at all energies, and the $R(s)$ values accounting for the contributions from both the continuum hadron production and the decays for $\psi(3770) \rightarrow \text{hadrons}$, respectively. All of these are corrected for the initial state radiative and vacuum polarization corrections. To extract the mass and widths of the $\psi(3770)$, we here, in the Letter, ana-




FIG. 1: The $R_{\text{uds}(c)+\psi(3770)}(s)$ versus the c.m. energy (see text).

lyze the quantity $R_{\text{uds}(c)+\psi(3770)}$ [9]. Table I summarizes the $R_{\text{uds}(c)+\psi(3770)}$ values reported in Ref. [9].

The determination of $M_{\psi(3770)}$, $\Gamma_{\psi(3770)}^{\text{tot}}$ and $\Gamma_{\psi(3770)}^{ee}$ is accomplished by simultaneously fitting the measured $R_{\text{uds}(c)+\psi(3770)}$ values listed in table I to the function that describes the combined $\psi(3770)$ resonance shape and non-resonant hadronic background. Assuming that there are no other new structures and effects, we use a pure p-wave zero order Breit-Wigner function with energy-dependent total widths to describe the $\psi(3770)$ production and its decay to inclusive hadrons. The $\psi(3770)$ resonance shape is taken as

$$R_{\psi(3770)}(s) = \frac{3s}{4\pi\alpha^2(0)} \frac{12\pi\Gamma_{\psi(3770)}^{ee}\Gamma_{\psi(3770)}(s)}{(s - M_{\psi(3770)}^2)^2 + [M_{\psi(3770)}\Gamma_{\psi(3770)}(s)]^2}, \quad (1)$$

with $\Gamma_{\psi(3770)}^{ee} = |1 - \Pi(s(1-x))|^2\Gamma_{\psi(3770)}^{ee}$, where $\Gamma_{\psi(3770)}^{ee}$ and $\Gamma_{\psi(3770)}^{ee}$ are the bare leptonic width excluding the vacuum polarization effects and experimental leptonic width including the vacuum polarization effects, respectively; $1/|1 - \Pi(s(1-x))|^2$ is the vacuum polarization correction function [10] including the contributions from all 1^{--} resonances, the QED continuum hadron spectrum as well as the contributions from the lepton pairs (e^+e^- , $\mu^+\mu^-$ and $\tau^+\tau^-$) [11]; x is a parameter related to the total energy of the emitted photons; the total width $\Gamma_{\psi(3770)}(s)$ is chosen to be energy dependent defined as

$$\Gamma_{\psi(3770)}(s) = \Gamma_{D^0\bar{D}^0}(s) + \Gamma_{D^+D^-}(s) + \Gamma_{\text{non-}D\bar{D}}(s), \quad (2)$$

in which [12]

$$\Gamma_{D^0\bar{D}^0}(s) = \Gamma_{\psi(3770)}^{\text{tot}}\theta_{00}\frac{(p_{D^0})^3}{(p_{D^0}^0)^3}\frac{1 + (rp_{D^0}^0)^2}{1 + (rp_{D^0}^0)^2}B_{00}, \quad (3)$$

$$\Gamma_{D^+D^-}(s) = \Gamma_{\psi(3770)}^{\text{tot}}\theta_{+-}\frac{(p_{D^+})^3}{(p_{D^+}^0)^3}\frac{1 + (rp_{D^+}^0)^2}{1 + (rp_{D^+}^0)^2}B_{+-} \quad (4)$$

TABLE I: Summary of the $R_{\text{uds}(c)+\psi(3770)}(s)$ values measured at 68 energy points.

$\sqrt{s}$ (GeV)	$R_{\text{uds}(c)+\psi(3770)}$	$\sqrt{s}$ (GeV)	$R_{\text{uds}(c)+\psi(3770)}$	$\sqrt{s}$ (GeV)	$R_{\text{uds}(c)+\psi(3770)}$	$\sqrt{s}$ (GeV)	$R_{\text{uds}(c)+\psi(3770)}$
3.6500	$2.157 \pm 0.035 \pm 0.086$	3.7584	$3.025 \pm 0.108 \pm 0.148$	3.7726	$3.777 \pm 0.145 \pm 0.185$	3.7826	$3.326 \pm 0.115 \pm 0.163$
3.6600	$2.131 \pm 0.105 \pm 0.085$	3.7596	$3.076 \pm 0.102 \pm 0.151$	3.7730	$3.563 \pm 0.120 \pm 0.175$	3.7838	$3.154 \pm 0.114 \pm 0.155$
3.6920	$2.034 \pm 0.092 \pm 0.081$	3.7608	$3.138 \pm 0.089 \pm 0.154$	3.7742	$3.373 \pm 0.113 \pm 0.165$	3.7850	$2.879 \pm 0.107 \pm 0.141$
3.7000	$2.079 \pm 0.079 \pm 0.083$	3.7620	$2.992 \pm 0.110 \pm 0.147$	3.7754	$3.641 \pm 0.125 \pm 0.178$	3.7862	$2.902 \pm 0.105 \pm 0.142$
3.7080	$2.197 \pm 0.083 \pm 0.088$	3.7622	$3.207 \pm 0.114 \pm 0.157$	3.7766	$3.498 \pm 0.119 \pm 0.171$	3.7874	$2.957 \pm 0.111 \pm 0.145$
3.7160	$2.177 \pm 0.086 \pm 0.087$	3.7634	$3.345 \pm 0.122 \pm 0.164$	3.7778	$3.570 \pm 0.121 \pm 0.175$	3.7886	$2.571 \pm 0.097 \pm 0.126$
3.7240	$2.125 \pm 0.086 \pm 0.085$	3.7646	$3.585 \pm 0.126 \pm 0.176$	3.7790	$3.360 \pm 0.117 \pm 0.165$	3.7898	$2.576 \pm 0.099 \pm 0.126$
3.7320	$2.156 \pm 0.086 \pm 0.086$	3.7658	$3.381 \pm 0.119 \pm 0.166$	3.7798	$3.477 \pm 0.136 \pm 0.170$	3.7900	$2.849 \pm 0.106 \pm 0.140$
3.7400	$2.190 \pm 0.099 \pm 0.088$	3.7670	$3.760 \pm 0.130 \pm 0.184$	3.7802	$3.427 \pm 0.125 \pm 0.168$	3.7950	$2.751 \pm 0.101 \pm 0.135$
3.7480	$2.371 \pm 0.106 \pm 0.116$	3.7682	$3.451 \pm 0.124 \pm 0.169$	3.7804	$3.382 \pm 0.137 \pm 0.166$	3.8000	$2.212 \pm 0.091 \pm 0.108$
3.7500	$2.517 \pm 0.085 \pm 0.123$	3.7694	$3.611 \pm 0.125 \pm 0.177$	3.7808	$3.336 \pm 0.129 \pm 0.163$	3.8100	$2.171 \pm 0.092 \pm 0.087$
3.7512	$2.637 \pm 0.090 \pm 0.129$	3.7706	$3.580 \pm 0.123 \pm 0.175$	3.7810	$3.464 \pm 0.138 \pm 0.170$	3.8200	$2.367 \pm 0.109 \pm 0.095$
3.7524	$2.615 \pm 0.095 \pm 0.128$	3.7714	$3.538 \pm 0.139 \pm 0.173$	3.7812	$3.396 \pm 0.130 \pm 0.166$	3.8300	$2.354 \pm 0.101 \pm 0.094$
3.7536	$2.652 \pm 0.093 \pm 0.130$	3.7716	$3.634 \pm 0.146 \pm 0.178$	3.7814	$3.514 \pm 0.124 \pm 0.172$	3.8400	$2.296 \pm 0.104 \pm 0.092$
3.7548	$2.733 \pm 0.093 \pm 0.134$	3.7718	$3.939 \pm 0.133 \pm 0.193$	3.7816	$2.944 \pm 0.137 \pm 0.144$	3.8500	$2.372 \pm 0.115 \pm 0.095$
3.7560	$2.585 \pm 0.090 \pm 0.127$	3.7720	$3.636 \pm 0.134 \pm 0.178$	3.7818	$3.140 \pm 0.125 \pm 0.154$	3.8600	$2.371 \pm 0.105 \pm 0.095$
3.7572	$2.942 \pm 0.107 \pm 0.144$	3.7722	$3.652 \pm 0.143 \pm 0.179$	3.7822	$3.253 \pm 0.124 \pm 0.159$	3.8720	$2.308 \pm 0.117 \pm 0.092$

and

$$\Gamma_{\text{non-}D\bar{D}}(s) = \Gamma_{\psi(3770)}^{\text{tot}} [1 - B_{00} - B_{+-}], \quad (5)$$

where p_D^0 and p_D are the momenta of the D mesons produced at the peak of the $\psi(3770)$ and at the c.m. energy $\sqrt{s}$, respectively; $\Gamma_{\psi(3770)}^{\text{tot}}$ is the total width of the $\psi(3770)$ at its peak, B_{00} and B_{+-} are the branching fractions for $\psi(3770) \rightarrow D^0\bar{D}^0$ and $\psi(3770) \rightarrow D^+D^-$, respectively, r is the interaction radius of the $c\bar{c}$, θ_{00} and θ_{+-} are the step functions to account for the thresholds of the $D^0\bar{D}^0$ and D^+D^- production, respectively. In the fit we take $\Gamma_{\psi(3770)}^{\text{tot}}$ and r as free parameters. Since we do not select the $D\bar{D}$ events from these data samples, we can not leave B_{00} and B_{+-} free in the fit. Instead we set the ratio $B_{00}/B_{+-} = 1.3 \pm 0.1$ [12, 13] and the branching fraction for $\psi(3770) \rightarrow D\bar{D}$ to be $B[\psi(3770) \rightarrow D\bar{D}] = B_{00} + B_{+-} = (84 \pm 7)\%$ [12, 14] in the fit.

The non-resonant background shape is taken as

$$R_{\text{uds}(c)}(s) = R_{\text{uds}}(s) + R_{(c)}(s), \quad (6)$$

with

$$R_{(c)}(s) = f_{D\bar{D}} \left[\left(\frac{p_{D^0}}{E_{D^0}} \right)^3 \theta_{00} + \left(\frac{p_{D^+}}{E_{D^+}} \right)^3 \theta_{+-} \right], \quad (7)$$

where $R_{\text{uds}}(s)$ is the R value for the continuum light hadron production; $R_{(c)}(s)$ is the R value due to the continuum $D\bar{D}$ ($D^0\bar{D}^0$ and D^+D^-) production; E_{D^0} and E_{D^+} are the energies of the D^0 and the D^+ mesons produced at $\sqrt{s}$, respectively; $f_{D\bar{D}}$ is a parameter to be fitted.

The expected $R_{\text{uds}(c)+\psi(3770)}(s)$ value is written as

$$R_{\text{uds}(c)+\psi(3770)}^{\text{expect}}(s) = \int_0^\infty ds'' G(s, s'') R_{\psi(3770)}(s) + R_{\text{uds}(c)}(s), \quad (8)$$

in which $G(s, s'')$ is the Gaussian function to describe the c.m. energy distribution of the BEPC machine [12]. In the fit we leave $R_{\text{uds}}(s)$ free, assuming that it is independent of the energy.

Figure 1 shows the measured $R_{\text{uds}(c)+\psi(3770)}(s)$ values with the fit, where the point with errors represents the $R_{\text{uds}(c)+\psi(3770)}(s)$ value, the curve (blue line) gives the fit to the data, the straight (yellow) line shows the quantity R_{uds} obtained from the fit and the dashed (red) line shows the variation of the quantity $R_{\text{uds}(c)}(s)$ with the c.m. energy as given in Eqs. (6-7). The χ^2/ndf for this fit is $94/61 = 1.5$. The larger χ^2 s are mainly from the four energy points, which are at $\sqrt{s} = 3.7886, 3.7898, 3.80$ and 3.81 GeV. The sum over the χ^2 s from these four energy points is 31. If we exclude these four points in the fit, we would obtain $\chi^2/\text{ndf} = 65/57 = 1.1$. The "dip" of the $R_{\text{uds}(c)+\psi(3770)}(s)$ values around 3.80 GeV, which obviously turn aside from the expected ones, may be due to some unknown effects [19]. Table II summarizes the results from the fit, where the first error is the statistical and the second systematic. The systematic errors on $\Gamma_{\psi(3770)}^{\text{tot}}$ and $\Gamma_{\psi(3770)}^{ee}$ arise mainly from the common systematic uncertainty on the measured quantity $R_{\text{uds}(c)+\psi(3770)}(s)$, while the systematic error on $M_{\psi(3770)}$ arises from the uncertainty in the energy calibration [12] of the BEPC collider. As a comparison, the same quantities measured by the BES Collaboration obtained by analyzing different cross section scan data sets [12] are

TABLE II: The measured mass $M_{\psi(3770)}$, total width $\Gamma_{\psi(3770)}^{\text{tot}}$ and leptonic width $\Gamma_{\psi(3770)}^{ee}$ of the $\psi(3770)$ along with those measured previously by the BES Collaboration [12].

$M_{\psi(3770)}$ (MeV)	$\Gamma_{\psi(3770)}^{\text{tot}}$ (MeV)	$\Gamma_{\psi(3770)}^{ee}$ (eV)	Note
$3772.4 \pm 0.4 \pm 0.3$	$28.6 \pm 1.2 \pm 0.2$	$279 \pm 11 \pm 13$	This work
$3772.2 \pm 0.7 \pm 0.3$	$26.9 \pm 2.4 \pm 0.3$	$251 \pm 26 \pm 11$	[12]

TABLE III: The leptonic branching fraction of the $\psi(3770)$.

Experiment	$B[\psi(3770) \rightarrow e^+e^-] \times 10^{-5}$
This work	$0.98 \pm 0.04 \pm 0.04$
BES [12]	$0.93 \pm 0.06 \pm 0.03$
PDG [8]	1.05 ± 0.14

also listed in the table. The fit also gives

$$R_{\text{uds}} = 2.121 \pm 0.023 \pm 0.084$$

in the energy region from 3.650 to 3.872 GeV, where the errors are statistical and systematic, respectively. The systematic uncertainty arises mainly from the uncertainty ($\sim 3.9\%$) in the measurements of $R_{\text{uds}(c)+\psi(3770)}(s)$. The measured R_{uds} value from this fit is consistent with the $R_{\text{uds}} = 2.141 \pm 0.025 \pm 0.085$ obtained by weighting the first 8 $R_{\text{uds}(c)+\psi(3770)}(s)$ values below the $D\bar{D}$ threshold [9]. In estimating the systematic uncertainties, the effects of the uncertainty in the measurement of the branching fraction for $\psi(3770) \rightarrow D\bar{D}$ on the measured quantities are also taken into account. The measured widths of the $\psi(3770)$ yield its leptonic branching fraction shown in table III.

Figure 2 illustrates the comparisons of the measured quantities (circle with error bar) with those measured by the MARK-I, DELCO and MARK-II Collaborations [16, 17, 18] from analyzing the cross section scan data. As a comparison we also plot the mass (dots) of the 1^3D_1 state of the $c\bar{c}$ system and the partial width (squares [3, 4]) for $\psi(3770) \rightarrow D\bar{D}$ predicted by the models [2, 3, 15] in recent years and calculated by the LQCD theory [5] in the figure.

With the mass, total width and leptonic width of the $\psi(3770)$, we extract the $\psi(3770)$ production cross section $\sigma_{\psi(3770)}^{\text{prd}}$ at $\sqrt{s} = M_{\psi(3770)}$, which excludes the initial state radiative corrections, and extract its corresponding observed cross section $\sigma_{\psi(3770)}^{\text{obs}}$, which includes the initial state radiative and vacuum polarization effects. Table IV summarizes the measured cross sections from this measurement and those measured by the MARK-II Collaboration and by the BES Collaboration from analysis of different cross section scan data [12].

In summary, we measured the mass of the $\psi(3770)$ with a precision of more than a factor of 4 better than the one of the PDG [8] average. The precise measurement of the mass combining with the measured masses of other $c\bar{c}$ states could be used to gauge the effect of systematic im-




FIG. 2: Comparison of the measured quantities obtained by analyzing the cross section scan data from different experiments along with those predicted by the models and by the LQCD theory; the dots are the predicted mass of the 1^3D_1 state of the $c\bar{c}$ system and the squares the expected partial width for $\psi(3770) \rightarrow D\bar{D}$ (see text).

TABLE IV: The cross sections for the $\psi(3770)$ production measured at $\sqrt{s} = M_{\psi(3770)}$, obtained by analyzing the cross section scan data over the $\psi(3770)$ or over both the $\psi(3770)$ and $\psi(3686)$.

Experiment	$\sigma^{\text{prd}}[e^+e^- \rightarrow \psi(3770)]$ [nb]	$\sigma^{\text{obs}}[e^+e^- \rightarrow \psi(3770)]$ [nb]
This work	$10.06 \pm 0.37 \pm 0.043$	$7.25 \pm 0.27 \pm 0.34$
BES [12]	$9.63 \pm 0.66 \pm 0.35$	$6.94 \pm 0.48 \pm 0.28$
MARK-II [18]		9.3 ± 1.4 [18]

provement on the calculation of the $c\bar{c}$ spectrum and to extract the $\alpha_s(s)$ at this mass scale [5]. Our measured $\Gamma_{\psi(3770)}^{\text{tot}} = 28.6 \pm 1.2 \pm 0.2$ MeV is significantly larger than the PDG average $\Gamma_{\psi(3770)}^{\text{tot}} = 23.0 \pm 2.7$ MeV [8], and with a precision more than a factor of 2 better than that of the PDG average. It is also obviously larger than the recently reasonable prediction by the coupled channel models [3, 4] by more than 7σ , where σ is the error of the measured value of the total width of $\psi(3770)$. Our measured leptonic width of the $\psi(3770)$ is consistent within error with the PDG average [8], but with a better accuracy than the PDG average. The measured leptonic branching fraction of the $\psi(3770)$ from this work is consistent within error with PDG average and with that measured by the BES Collaboration previously [12], but with a precision of more than a factor of 2 better than the PDG average. We measured the $\psi(3770)$ production cross section to be $\sigma_{\psi(3770)}^{\text{obs}} = 7.25 \pm 0.27 \pm 0.34$

nb at $\sqrt{s} = M_{\psi(3770)}$. These improved measurements of the quantities would be helpful for us to understand the nature of the $\psi(3770)$ resonance. From the analysis we also extracted $R_{uds} = 2.121 \pm 0.023 \pm 0.084$ in the energy range from 3.650 to 3.872 GeV. This information would be beneficial for us in the understanding of the continuum hadron production in or nearby the resonance region(s). The measured R_{uds} can directly be used to extract the $\alpha_s(s)$ at this energy scale.

The BES collaboration thanks the staff of BEPC for their hard efforts. This work is supported in part by the National Natural Science Foundation of China under contracts Nos. 19991480, 10225524, 10225525, the Chinese Academy of Sciences under contract No. KJ 95T-03, the 100 Talents Program of CAS under Contract Nos. U-11, U-24, U-25, and the Knowledge Innovation Project of CAS under Contract Nos. U-602, U-34(IHEP); by the National Natural Science Foundation of China under Contract No. 10175060(USTC), and No. 10225522(Tsinghua University).

-
- [1] E. Eichten, K. Gottfried *et al.*, Phys. Rev. Lett. **34** 369 (1975).
 - [2] E. Eichten, K. Gottfried *et al.*, Phys. Rev. D **21** 203 (1980).
 - [3] K. Heikkilä and N.A. Törnqvist, Phys. Rev. D **29** 110 (1983).
 - [4] E. Eichten, K. Lane, C. Quigg Phys. Rev. Lett. **69** 094019 (2004).
 - [5] C.T.H. Davies, K. Hornbostel *et al.*, Phys. Rev. D **52** 6519 (1995).
 - [6] M. Ablikim *et al.* (BES Collaboration), Phys. Lett. B **603**, 130 (2004); Nucl. Phys. B **727**, 395 (2005); Phys. Lett. B **641**, 145 (2006); Phys. Rev. Lett. **97**, 121801 (2006); HEP & NP **28(4)**, 325 (2004); Phys. Lett. B **605**, 63 (2005); Phys. Rev. D **70**, 077101 (2004); Phys. Rev. D **72**, 072007 (2005); G. Rong (for BES Collaboration), Int. J. Mod. Phys., **21**, 5416 (2006). G. Rong, D.H. Zhang and J.C. Chen, hep-ex/0506051.
 - [7] N. E. Adam *et al.* (CLEO Collaboration), Phys. Rev. Lett. **96**, 082004 (2006); T. E. Coans *et al.* (CLEO Collaboration), Phys. Rev. Lett. **96**, 182002 (2006); G. S. Huang *et al.* (CLEO Collaboration), Phys. Rev. Lett. **96**, 032003 (2006); G. S. Adams *et al.* (CLEO Collaboration), Phys. Rev. D **73**, 012002 (2006); D. Cronin-Hennessy *et al.* (CLEO Collaboration), Phys. Rev. D **74**, 012005 (2006); R. A. Briere *et al.* (CLEO Collaboration), Phys. Rev. D **74**, 031106 (2006); D. Beson *et al.* (CLEO Collaboration), Phys. Rev. Lett. **96**, 092002 (2006).
 - [8] W-M Yao *et al.* (Particle Data Group), J. Phys. G **33**, 1 (2006).
 - [9] M. Ablikim *et al.* (BES Collaboration), hep-ex/0612054; to be published in Phys. Rev. Lett. (in press).
 - [10] F.A. Berends and G.J. Komen, Phys. Lett. B **63**, 432 (1976); Andrej B. Arbuzov, Eduard A. Kuraev *et al.*, JHEP10, 006 (1997).
 - [11] D.H. Zhang, G. Rong and C.J. Chen, Phys. Rev. D **74** 054012 (2006).
 - [12] M. Ablikim *et al.* (BES Collaboration), Phys. Rev. Lett. **97** 121801 (2006).
 - [13] M. Ablikim *et al.* (BES Collaboration), Nucl. Phys. B **727** 395 (2005).
 - [14] M. Ablikim *et al.* (BES Collaboration), Phys. Lett. B **641** 145 (2006).
 - [15] S. Ono, Phys. Rev. D **20**, 2975(1979); S. Ono, Z. Phys. C **8**, 7 (1981); A. Martin, Phys. Lett. B **100**, 511 (1981); L. Richardson, Phys. Lett. B **82**, 272 (1979); W. Buchmüller and S.-H. H. Tye, Phys. Rev. D **24**, 132 (1981); G. Bhanot and S. Rudaz, Phys. Lett. B **78**, 119 (1978); H. Krasemann and S. Ono, Nucl. Phys. B **154**, 283 (1979).
 - [16] P.A. Rapidis *et al.* (MARK-I Collaboration), Phys. Rev. Lett. **39** (1977) 526.
 - [17] W. Bacino *et al.* (DELCO Collaboration), Phys. Rev. Lett. **40** (1978) 671.
 - [18] R.H. Schindler *et al.* (MARK-II Collaboration), Phys. Rev. D **21** (1980) 2716.
 - [19] The analysis author would like to suggest the BES-III (BES-III at BEPC-II) and/or CLEO (CLEO-c at CERN) Collaborations to check whether this "dip" is due to physical reason with larger statistical cross section scan data to be collected at more sampling points around $\sqrt{s} = 3.80$ GeV in the future.